Exact Model Reduction for Discrete-Time Conditional Quantum Dynamics

Tommaso Grigoletto and Francesco Ticozzi

Milan, December 16th, 2024



What are the minimal resources needed to reproduce a target quantum process?

Finding the minimal resources allows us to reduce the model's description.

Why is quantum model reduction interesting?

- Efficient quantum simulation (on classical and quantum computers);
- Efficient implementations of:
 - controllers,
 - error suppression schemes,
 - quantum filters;
- Easier models to study;
- Proving optimality of quantum algorithms;
- Probing "quantumness" of processes;
- Efficient generation of quantum trajectories (Monte Carlo methods).

Dynamics of interest: Conditional dynamics

- CPTP evolution followed by generalized measurement;
- Imperfect measurement;
- Dynamics conditioned on the measurement outcome.

In general we assume to have a set of CP $\{\mathcal{M}_k\}_{k\in\Omega}$, s.t $\sum_{k\in\Omega}\mathcal{M}_k^{\dagger}(\mathbf{1})=\mathbf{1}$ with Ω the set of possible outcomes.

If the state of the system is ρ , the outcome $k \in \Omega$ is observed with probability

$$\mathbb{P}_{\rho}[\mathcal{M}=k] = \operatorname{tr}[\mathcal{M}_k(\rho)].$$

The state of the system, conditioned on the measurement outcome, then is

$$\rho|_{\mathcal{M}=k} = \frac{\mathcal{M}_k(\rho)}{\operatorname{tr}[\mathcal{M}_k(\rho)]}.$$

Linear dynamics

The conditioning rule $\rho|_{\mathcal{M}=k} = \frac{\mathcal{M}_k(\rho)}{\operatorname{tr}[\mathcal{M}_k(\rho)]}$ is non-linear.

But we can consider **un-normalized states** $\tilde{\rho}$ so that the dynamics becomes

$$\tilde{\rho}(t+1) = \mathcal{M}_{k_{t+1}}[\tilde{\rho}(t)]$$
.

The probability of a sequence of outcomes then is:

$$\mathbb{P}[\mathcal{M}_{0:t} = k_{0:t}] = \operatorname{tr}[\mathcal{M}_{k_{0:t}}(\rho_0)] = \operatorname{tr}[\mathcal{M}_{k_t} \circ \cdots \circ \mathcal{M}_{k_1} \circ \mathcal{M}_{k_0}(\rho_0)]$$

To compute the normalized state or expectation values one can simply re-normalize:

$$\rho(t) = \frac{\tilde{\rho}(t)}{\mathbb{P}[\mathcal{M}_{0:t} = k_{0:t}]} \qquad \langle O(t) \rangle = \text{tr}[O\rho(t)] = \frac{\text{tr}[O\tilde{\rho}(t)]}{\mathbb{P}[\mathcal{M}_{0:t} = k_{0:t}]}$$

Quantities of interest:

In many practical settings, one is not interested in the entire state ρ .

We thus assume to only be interest in reproducing the expectation value of a **set of observables of interest** $\{O_j\}$ that include the identity, $\mathbf{1} \in \{O_j\}$.


We can compactly represent these as a linear output map $\mathcal{C}:\mathfrak{B}(\mathcal{H})\to\mathscr{Y}$: $\mathcal{C}[\rho]=\sum_j E_j\mathrm{tr}[O_j\rho].$

In the following we focus in the **conditional dynamics (CD)** ($\{\mathcal{M}_k\}, \mathcal{C}$):

$$\begin{cases} \tilde{\rho}(t+1) &= \mathcal{M}_{k_{t+1}}[\tilde{\rho}(t)] \\ Y(t) &= \mathcal{C}[\tilde{\rho}(t)] \end{cases}$$

Similar to Hidden Markov Models.

Monitored spin chain - A toy model -



$$\mathcal{E}(\cdot) = e^{-iH} \cdot e^{iH} \qquad \text{where} \qquad H = \delta \sum_{i=1}^{N-1} \sigma_x^{(j)} \sigma_x^{(j+1)}$$

We assume to perform a projective on the last spin with probability $p \in (0,1)$.

$$\mathcal{K}_{-1}(\cdot) = p\mathbf{1} \cdot \mathbf{1}, \qquad \mathcal{K}_{0}(\cdot) = (1-p)\Pi_{0} \cdot \Pi_{0}, \qquad \mathcal{K}_{1}(\cdot) = (1-p)\Pi_{1} \cdot \Pi_{1}$$

where Π_0, Π_1 are the eigenprojectors of $\sigma_z^{(N)}$.

$$\mathcal{M}_k = \mathcal{K}_k \circ \mathcal{E}$$

We are actually interested in reproducing the evolution of the first spin, i.e.

$$\rho_1(t) = \operatorname{tr}_{\bar{1}}[\rho(t)]$$
 $\mathcal{C}(\cdot) = \operatorname{tr}_{\bar{1}}(\cdot)$

The problem: Quantum model reduction

Given a CD $(\{\mathcal{M}_k\}, \mathcal{C})$ find:

- another CD $(\{\check{\mathcal{M}}_k\},\check{\mathcal{C}});$
- a linear map $\Phi: \mathbb{C}^{n\times n}\to \mathbb{C}^{r\times r}$ so that $\check{\rho}_0=\Phi[\rho_0]$ such that

$$\mathcal{C}[\mathcal{M}_{k_{0:t}}(\rho_0)] = \check{\mathcal{C}}[\check{\mathcal{M}}_{k_{0:t}}(\check{\rho}_0)]$$

for all $t \geq 0$ all $\rho_0 \in \mathfrak{D}(\mathcal{H})$, and for all sequences of outcomes $k_{0:t}$.

Observable space

Let us define

$$\mathscr{O} = \operatorname{span}\{O_j, \mathcal{M}_{k_{0:t}}^{\dagger}(O_j), \quad \forall j, \quad \forall k_{0:t}, \quad \forall t\}$$

the (Krylov) operator space that contains the <u>observables of interest evolved in</u> Heisenberg picture, for all possible trajectories.

One can see it as the orthogonal to the "usual" non-observable subspace \mathcal{N} .

FACT: \mathscr{O} is the smallest \mathcal{M}_k^{\dagger} -invariant operator space that contains $\{O_j\}$.

 \mathscr{O} contains all the degrees of freedom we need to reproduce $\mathscr{C}[\mathcal{M}_{k_0:t}(\rho_0)]$.

Problem:

Restricting the model onto \mathcal{O} does not ensure Complete Positivity

How do we ensure that physical (probability) constraints are satisfied?

The solution: *-algebras

We define a *-algebra $\mathscr A$ as an operator space closed under matrix multiplication and adjoint action.

$$X,Y\in\mathscr{A}\quad\Rightarrow\quad X+Y\in\mathscr{A}\qquad X^\dagger,Y^\dagger\in\mathscr{A}\qquad\text{and}\qquad XY\in\mathscr{A}$$

It is the fundamental mathematical structure that supports a **quantum probability space**.

Conditional expectations

A conditional expectation $\mathbb{E}_{\mathscr{A},\rho}[\cdot]$ is a CP unital proj. onto a *-algebra \mathscr{A} (and such that $\operatorname{tr}[\rho \mathbb{E}_{\mathscr{A},\rho}[X]] = \operatorname{tr}[\rho X]$ for all $X \in \mathcal{B}(\mathcal{H})$).


When $1 \in \mathcal{A}$, $\mathbb{E}_{\mathcal{A},1/n}[\cdot]$ exists and is an orthogonal proj. hence CPTP.

 $\mathbb{E}_{\mathscr{A},1/n}[\cdot]$ can be factorized in two non-square CPTP isometries

$$\mathbb{E}_{\mathscr{A},\mathbf{1}/n}[\cdot] = \mathcal{J}\mathcal{R}.$$

Given a CPTP map A, its reduction onto the algebra A,

$$\mathcal{A}|_{\mathscr{A}} = \mathcal{R}\mathcal{A}\mathcal{J}$$



Observable CPTP reduction

We leverage the knowledge of the quantities of interest to reduce the model:

- 1. Find the observable subspace \mathscr{O} .
- **2**. Close its orthogonal complement to a *-algebra $\mathscr{A} = alg(\mathscr{O})$;
- **3**. Compute the **cond. exp.** $\mathbb{E}_{\mathscr{A},1/n}[\cdot]$ and its CPTP factors \mathcal{R} and \mathcal{J} ;
- 4. Use the two factors to reduce the dynamics:

$$\check{\mathcal{M}}_k = \mathcal{R} \mathcal{M}_k \mathcal{J}, \qquad \check{\mathcal{C}} = \mathcal{C} \mathcal{J}, \qquad \Phi = \mathcal{R}.$$

Note that $\check{\mathcal{M}}_k$ are CP and also $\sum_{k\in\Omega}\check{\mathcal{M}}_k^\dagger(1)=1$ as $\mathcal{R}^\dagger,\mathcal{J}^\dagger$ are unital.

Reduction of Measurements and Dynamics

In many cases we have $\mathcal{M}_k = \mathcal{K}_k \circ \mathcal{E}$ with \mathcal{E} a CPTP dynamics and $\mathcal{K}_k(\cdot) = M_k \cdot M_k^{\dagger}$ a generalized measurement, $\sum_{k \in \Omega} M_k^{\dagger} M_k = \mathbf{1}$.


Assuming one of the following:

- 1. $\exists \{\lambda_k\}$ s.t. $\sum_k \mathcal{M}_k = \mathcal{E}$;
- 2. \mathcal{N}^{\perp} is \mathcal{E}^{\dagger} -invariant;
- 3. \mathscr{A} is \mathcal{K}_k -invariant $\forall k$;
- **4.** \mathscr{A} is \mathcal{E}^{\dagger} -invariant;

We have

$$\check{\mathcal{M}}_k = \mathcal{R}\mathcal{M}_k\mathcal{J} = \mathcal{R}\mathcal{K}_k\mathcal{J} \ \mathcal{R}\mathcal{E}\mathcal{J} = \check{\mathcal{K}}_k\check{\mathcal{E}}$$

Monitored spin chain - A toy model -



$$\mathcal{E}(\cdot) = e^{-iH} \cdot e^{iH} \qquad \text{where} \qquad H = \delta \sum_{i=1}^N \sigma_x^{(j)} \sigma_x^{(j)}$$

We assume to perform a projective on the last spin with probability $p \in (0,1)$.

$$\mathcal{K}_{-1}(\cdot) = p\mathbf{1} \cdot \mathbf{1}, \qquad \mathcal{K}_{0}(\cdot) = (1-p)\Pi_{0} \cdot \Pi_{0}, \qquad \mathcal{K}_{1}(\cdot) = (1-p)\Pi_{1} \cdot \Pi_{1}$$

where Π_0, Π_1 are the eigenprojectors of $\sigma_z^{(N)}$.

$$\mathcal{M}_k = \mathcal{K}_k \circ \mathcal{E}$$

We are actually interested in reproducing the evolution of the first spin, i.e.

$$\rho_1(t) = \operatorname{tr}_{\bar{1}}[\rho(t)]$$
 $\mathcal{C}(\cdot) = \operatorname{tr}_{\bar{1}}(\cdot)$

Monitored spin chain - Spaces

For $N \ge 4$, regardeless of N we have $\dim(\mathcal{O}) = 18$. The evolution of $\rho_1(t)$ is only influences by spins 1, 2, N - 1, N.

Moreover,

$$\mathscr{A} \simeq \mathbb{C}^{4 \times 4} \oplus \mathbb{C}^{4 \times 4}.$$

and is \mathcal{K}_k -invariant.

The reduced model is thus a classical mixture of two two-qubit systems and $\check{\mathcal{M}}_k = \check{\mathcal{K}}_k \check{\mathcal{E}}$.

$$\xi(t) = \left[\begin{array}{c|c} \xi_0(t) & 0 \\ \hline 0 & \xi_1(t) \end{array} \right] \in \mathbb{C}^{8 \times 8}$$

Take home ideas

1. We need algebras to define a quantum probability space;

2. **Conditional expectations** provide CPTP reduction.

Conclusion



- Framework for model reduction of statistical dynamics (CPTP)
- Applied to discrete-time quantum trajectories [arXiv:2403.12575];



- The framework is very general and has been applied to:
 - (deterministic) Discrete-time case [arXiv:2307.06319];
 - (deterministic) Continuous-time case NEW! (joint with LV) [arXiv:2412.05102];
 - Continuous-time quantum trajectories (in preparation).
- Outlook
 - Approximate model reduction;
 - Connection with adiabatic elimination techniques;

Thanks for your attention!



Our group's website →